## **MOTRIN**®

# Dosing for Infants and Children

Concentrated Motrin Infants Drops



From Your Healthcare Professional

DOSE: Every 6-8 hours as needed. DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.

If possible, use weight to dose; otherwise use age.

> mL = milliliter tsp = teaspoonful



#### Infants' MOTRIN® **Concentrated Drops**

Active Ingredient: Ibuprofen 50 mg (NSAID)1 (in each 1.25 mL)

\*Nonsteroidal anti-inflammatory drug

Available in:



#### Children's MOTRIN® **Oral Suspension**

Active Ingredient:

Ibuprofen 100 mg (NSAID)1 (in each 5 mL or 1 tsp)

Nonsteroidal anti-inflammatory drug Available in:

WEIGHT	AGE	Original Berry Haver Use only as directed.  dye- Dye-free free Berry	Use only as directed.  dye Dye-free Berry
6-11 lbs	0-5 mos	Do not use	
12-17 lbs	6-11 mos	1.25 mL	-
18-23 lbs	12-23 mos	1.875 mL	
24-35 lbs	2-3 yrs	<del>-</del>	5 mL (1 tsp)
36-47 lbs	4-5 yrs		7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs		10 mL (2 tsp)
60-71 lbs	9-10 yrs		12.5 mL (2½ tsp)
72-95 lbs	11 yrs	<del></del>	15 mL (3 tsp)

### IMPORTANT INSTRUCTIONS for Proper Use

Today's Date: This dosing recommendation from your doctor will expire in 14 DAYS.

- Read and follow the label on all MOTRIN® products
- Take every 6-8 hours as needed. Do NOT exceed more than 4 doses in 24 hours
- Do NOT administer longer than 10 days, unless directed by a doctor
- · Ask a doctor or pharmacist before use if the child is taking any other drug containing an NSAID† (prescription or non-prescription)
- MOTRIN® contains ibuprofen
- Use only the dosing device that comes with a specific product

Nonsteroidal anti-inflammatory drug



#### Kids' Wellness Tracker

Free, all-in-one tool from the makers of Children's TYLENOL® and Children's MOTRIN®. Makes it easy to track your child's height, weight, vaccines, symptoms, and medicines plus calculate dosing.



App Store Download now at the App Store App Store is a service mark of Apple Inc. Standard download rates may apply